

On Decomposition of Continuity Via $*\mu$ -Closed Sets

Selvaraj Ganesan

Assistant Professor, PG & Research Department of Mathematics, Raja Doraisingam Government Arts College,
Sivagangai-630561, Tamil Nadu, India.

(Affiliated to Alagappa University, Karaikudi, Tamil Nadu, India)

sgsgsgsgsg77@gmail.com

(Received: January / 2020. Accepted: April / 2020.)

Abstract: There are various types of generalization of continuous functions in the development of topology. Recently some decompositions of continuity are obtained by various authors with the help of generalized continuous functions in topological spaces. In this paper we obtain a decomposition of continuity using a generalized continuity called $*\mu$ -continuity in topology.

Keywords: $*\mu$ -closed set, $*\mu$ lc*-set, $*\mu$ -continuous function, $*\mu$ lc*-continuous function. **2010 Mathematics**

Subject Classification: 54C10, 54C08, 54C05.

1 INTRODUCTION

Different types of generalizations of continuous function were introduced and studied by various authors in the recent development of topology. The decomposition of continuity is one of the many problems in general topology. Tong [13] introduced the notions of A-sets and A-continuity and established a decomposition of continuity. Also Tong [14] introduced the notions of B-sets and B-continuity and used them to obtain another decomposition of continuity and Ganster and Reilly [8] improved Tong's decomposition result. Recently, various decompositions of continuity have been established, for example [2, 3, 4, 5, 6, 7, 8, 11]. In this paper, we obtain a decomposition of continuity in topological spaces by using $*\mu$ -continuity.

2 PRELIMINARIES

Throughout this paper (X, τ) and (Y, σ) (or X and Y) represent topological spaces on which no separation axioms are assumed unless otherwise mentioned. For a subset A of a space (X, τ) , $cl(A)$ and $int(A)$ denote the closure of A and the interior of A respectively.

We recall the following definitions which are useful in the sequel.

Definition 2.1 A subset A of a space (X, τ) is called a semi-open set [9] if $A \subseteq cl(int(A))$.

The complements of semi-open set is said to be semi-closed.

The semi-closure [1] of a subset A of X , denoted by $scl(A)$ is defined to be the intersection of all semi-closed sets of (X, τ) containing A . It is known that $scl(A)$ is a semi-closed set.

Definition 2.2 A subset A of a space (X, τ) is called:



1. a \hat{g} -closed set [17] ($= \omega$ -closed set [12]) if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi-open in (X, τ) . The complement of \hat{g} -closed set is called \hat{g} -open.
2. a *g -semi closed (briefly *gs -closed) set [10, 15] if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is \hat{g} -open in (X, τ) . The complement of *gs -closed set is called *gs -open.
3. a $^*\mu$ -closed set [16] if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is *gs -open in (X, τ) . The complement of $^*\mu$ -closed set is called $^*\mu$ -open.

Definition 2.3 [16] A function $f : (X, \tau) \rightarrow (Y, \sigma)$ is said to be $^*\mu$ -continuous if for each closed set V of Y , $f^{-1}(V)$ is $^*\mu$ -closed in X .

Proposition 2.4 [16] Every closed set is $^*\mu$ -closed but not conversely.

Proposition 2.5 [16] Every continuous function is $^*\mu$ -continuous but not conversely.

3 DECOMPOSITION OF CONTINUITY

In this section, we obtain a decomposition of continuity in topological spaces by using $^*\mu$ -continuity.

To obtain a decomposition of continuity, we first recall the notion of $^*\mu lc^*$ -continuous function in topological spaces by using $^*\mu$ -continuity and prove that a function is continuous if and only if it is both $^*\mu$ -continuous and $^*\mu lc^*$ -continuous.

We introduce the following definition.

Definition 3.1 A subset A of a space (X, τ) is called $^*\mu lc^*$ -set if $A = M \cap N$, where M is *gs -open and N is closed (X, τ) .

Example 3.2 Let $X = \{a, b, c\}$ with $\tau = \{\phi, \{a\}, X\}$. Then $\{c\}$ is $^*\mu lc^*$ -set in (X, τ) .

Remark 3.3 Every closed set is $^*\mu lc^*$ -set but not conversely.

Example 3.4 Let $X = \{a, b, c\}$ with $\tau = \{\phi, \{b\}, X\}$. Then $\{a, b\}$ is $^*\mu lc^*$ -set but not closed in (X, τ) .

Remark 3.5 $^*\mu$ -closed and $^*\mu lc^*$ -sets are independent of each other.

Example 3.6 Let $X = \{a, b, c\}$ with $\tau = \{\phi, \{a, b\}, X\}$. Then $\{a, c\}$ is an $^*\mu$ -closed set but not $^*\mu lc^*$ -set in (X, τ) . Also, $\{a, b\}$ is $^*\mu lc^*$ -set in (X, τ) but it is not $^*\mu$ -closed.

Proposition 3.7 Let (X, τ) be a topological space. Then a subset A of (X, τ) is closed if and only if it is both $^*\mu$ -closed and $^*\mu lc^*$ -set.

Proof Necessity is trivial. To prove the sufficiency, assume that A is both $^*\mu$ -closed and $^*\mu lc^*$ -set. Then $A = M \cap N$, where M is *gs -open and N is closed in (X, τ) . Therefore, $A \subseteq M$ and $A \subseteq N$ and so by hypothesis, $cl(A) \subseteq M$ and $cl(A) \subseteq N$. Thus $cl(A) \subseteq M \cap N = A$ and hence $cl(A) = A$ i.e., A is closed in (X, τ) .

Definition 3.8 A function $f : (X, \tau) \rightarrow (Y, \sigma)$ is said to be $^*\mu\text{lc}^*$ -continuous if for each closed set V of (Y, σ) , $f^{-1}(V)$ is $^*\mu\text{lc}^*$ -set in (X, τ) .

Example 3.9 Let $X = Y = \{a, b, c\}$ with $\tau = \{\phi, \{c\}, X\}$ and $\sigma = \{\phi, \{b\}, \{a, c\}, Y\}$. Let $f : (X, \tau) \rightarrow (Y, \sigma)$ be the identity function. Then f is $^*\mu\text{lc}^*$ -continuous function.

Remark 3.10 Every continuous function is $^*\mu\text{lc}^*$ -continuous but not conversely.

Remark 3.11 $^*\mu$ -continuity and $^*\mu\text{lc}^*$ -continuity are independent of each other.

Example 3.12 Let $X = Y = \{a, b, c\}$ with $\tau = \{\phi, \{a, b\}, X\}$ and $\sigma = \{\phi, \{b\}, Y\}$. Let $f : (X, \tau) \rightarrow (Y, \sigma)$ be the identity function. Then f is $^*\mu$ -continuous but not $^*\mu\text{lc}^*$ -continuous. Let $\sigma = \{\phi, \{c\}, Y\}$ and $g : (X, \tau) \rightarrow (Y, \sigma)$ be the identity function. Then g is $^*\mu\text{lc}^*$ -continuous but it is not $^*\mu$ -continuous.

We have the following decomposition for continuity.

Theorem 3.13 A function $f : (X, \tau) \rightarrow (Y, \sigma)$ is continuous if and only if it is both $^*\mu$ -continuous and $^*\mu\text{lc}^*$ -continuous.

Proof Assume that f is continuous. Then by Proposition 2.5 and Remark 3.10, f is both $^*\mu$ -continuous and $^*\mu\text{lc}^*$ -continuous.

Conversely, assume that f is both $^*\mu$ -continuous and $^*\mu\text{lc}^*$ -continuous. Let V be a closed subset of (Y, σ) . Then $f^{-1}(V)$ is both $^*\mu$ -closed and $^*\mu\text{lc}^*$ -set. By Proposition 3.7, $f^{-1}(V)$ is a closed set in (X, τ) and so f is continuous.

References

- [1] Crossley, S. G. and Hildebrand, S. K.: Semi-closure, Texas J. Sci, 22(1971), 99-112.
- [2] Ekici, E. and Jafari, S.: On D-sets, DS-sets and decompositions of continuous, A-continuous and AB-continuous functions, Italian Journal of Pure and Applied Mathematics., 24 (2008), 255-264.
- [3] Ekici, E., Jafari, S. and Latif, R. M.: On a finer topological space than τ_θ and some maps, Italian Journal of Pure and Applied Mathematics., N 27(2010), 293-304.
- [4] Ekici, E.: On C^* -sets and decompositions of continuous and $\eta\zeta$ -continuous functions, Acta Mathematica Hungarica., 117 (4) (2007), 325-333.
- [5] Ekici, E. and Noiri, T.: Decompositions of continuity, α -continuity and AB-continuity, Chaos, Solitons and Fractals., 41 (2009), 2055-2061.
- [6] Ekici, E.: On e-open sets, DP*-sets and DPE*-sets and decompositions of continuity, Arabian Journal for Science and Engineering., 33(2A) (2008), 269-282.
- [7] Ganesan, S.: Decomposition of continuity via μ -closed sets, Antarctica Journal of Mathematics., 10(3)(2013), 203-206.
- [8] Ganster, M. and Reilly, I. L.: A decomposition of continuity, Acta Mathematica Hungarica., 56(1990), 299-301.

- [9] Levine, N.: Semi-open sets and semi-continuity in topological spaces, *Amer. Math. Monthly*, 70(1963), 36-41.
- [10] Palaniappan, N., Antony Rex Rodrigo, J. and Missier, S. P.: On $\hat{\eta}^*$ -closed sets in topological spaces, *International Journal of General Topology.*, 1(1) (June 2008), 77-85.
- [11] Ravi, O., Ganesan, S. and Chandrasekar, S.: Decomposition of continuity via g^* -closed sets, *Italian Journal of Pure and Applied Mathematics.*, in press.
- [12] Sheik John, M.: A study on generalizations of closed sets and continuous maps in topological and bitopological spaces, Ph.D Thesis, Bharathiar University, Coimbatore, September 2002.
- [13] Tong, J.: A decomposition of continuity, *Acta Mathematica Hungarica.*, 48(1986), 11-15.
- [14] Tong, J.: On decomposition of continuity in topological spaces, *Acta Mathematica Hungarica.*, 54(1989), 51-55.
- [15] Veera Kumar, M. K. R. S.: μp -closed sets in topological spaces, *Antarctica J. Math.*, 2(1) (2005), 31-52.
- [16] Veera Kumar, M. K. R. S.: $^*\mu$ -closed sets, *Antarctica J. Math.*, 2(2) (2005), 223-237.
- [17] Veera Kumar, M. K. R. S.: \hat{g} -closed sets in topological spaces, *Bull. Allahabad Math. Soc.*, 18(2003), 99-112.