

A Generalized Summation Formula

Salahuddin¹, Vinti²

¹Department of Mathematics, PDM University, Bahadurgarh 124507, Haryana, India

²Department of Mathematics, PDM University, Bahadurgarh 124507, Haryana, India
vsludn@gmail.com

Abstract

In this paper, we have developed the generalized expression of

$${}_2F_1 \left[\begin{matrix} a, b & ; & 1 \\ \frac{a+b-n}{2} & ; & 2 \end{matrix} \right]$$

and it's corresponding integral form .

Key Words : Contiguous relation, Summation formulae.

2010 MSC NO: 33C05, 33C20, 33C60

1. Introduction

Appell's Hypergeometric Function $F_1(x, y), F_2(x, y), F_3(x, y), F_4(x, y)$ are defined as(see[6]):

$$F_1[a; b, c; d; x, y] = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(a)_{m+n}(b)_m(c)_n x^m y^n}{(d)_{m+n} m! n!}, \quad \max\{|x|, |y|\} < 1 \tag{1.1}$$

$$F_2[a; b, c, d, e; x, y] = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(a)_{m+n}(b)_m(c)_n x^m y^n}{(d)_m (e)_n m! n!}, \quad |x| + |y| < 1 \tag{1.2}$$

$$F_3[a, b; c, d; e; x, y] = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(a)_m (b)_n (c)_m (d)_n x^m y^n}{(e)_{m+n} m! n!}, \quad \max\{|x|, |y|\} < 1 \tag{1.3}$$

$$F_4[a, b; c, d; x, y] = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(a)_{m+n}(b)_{m+n} x^m y^n}{(c)_m (d)_n m! n!}, \quad \sqrt{|x|} + \sqrt{|y|} < 1 \tag{1.4}$$

A generalized hypergeometric function ${}_pF_q(a_1, \dots, a_p; b_1, \dots, b_q; z)$ is a function which can be defined in the form of a hypergeometric series, i.e., a series for which the ratio of successive terms can be written

$$\frac{c_{k+1}}{c_k} = \frac{P(k)}{Q(k)} = \frac{(k + a_1)(k + a_2) \dots (k + a_p)}{(k + b_1)(k + b_2) \dots (k + b_q)(k + 1)} z. \tag{1.5}$$

Where $k + 1$ in the denominator is present for historical reasons of notation[Koepf p.12(2.9)], and the resulting generalized hypergeometric function is written

$${}_pF_q \left[\begin{matrix} a_1, a_2, \dots, a_p & ; \\ b_1, b_2, \dots, b_q & ; \end{matrix} \right] z = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \dots (a_p)_k z^k}{(b_1)_k (b_2)_k \dots (b_q)_k k!} \tag{1.6}$$

where the parameters b_1, b_2, \dots, b_q are positive integers.

The ${}_pF_q$ series converges for all finite z if $p \leq q$, converges for $|z| < 1$ if $p = q + 1$, diverges for all $z, z \neq 0$ if $p > q + 1$ [Luke p.156(3)].



The function ${}_2F_1(a, b; c; z)$ corresponding to $p = 2, q = 1$, is the first hypergeometric function to be studied (and, in general, arises the most frequently in physical problems), and so is frequently known as "the" hypergeometric equation or, more explicitly, Gauss's hypergeometric function [Gauss p.123-162]. To confuse matters even more, the term "hypergeometric function" is less commonly used to mean closed form, and "hypergeometric series" is sometimes used to mean hypergeometric function.

The hypergeometric functions are solutions of Gaussian hypergeometric linear differential equation of second order

$$z(1 - z)y'' + [c - (a + b + 1)z]y' - aby = 0 \tag{1.7}$$

The solution of this equation is

$$y = A_0 \left[1 + \frac{ab}{1!c}z + \frac{a(a+1)b(b+1)}{2!c(c+1)}z^2 + \dots \right] \tag{1.8}$$

This is the so-called regular solution, denoted

$${}_2F_1(a, b; c; z) = \left[1 + \frac{ab}{1!c}z + \frac{a(a+1)b(b+1)}{2!c(c+1)}z^2 + \dots \right] = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k z^k}{(c)_k k!} \tag{1.9}$$

which converges if c is not a negative integer for all $|z| < 1$ and on the unit circle $|z| = 1$ if $R(c - a - b) > 0$.

It is known as Gauss hypergeometric function in terms of Pochhammer symbol $(a)_k$ or generalized factorial function.

Digamma function:

In mathematics, the digamma function is defined as the logarithmic derivative of the gamma function [Abramowitz p.258-259]

$$\psi(x) = \frac{d}{dx} \ln\{\Gamma(x)\} = \frac{\Gamma'(x)}{\Gamma(x)} \tag{1.10}$$

The digamma function is often denoted as $\psi^0(x)$.

Pochhammer symbol:

In mathematics, the falling factorial or Pochhammer symbol (sometimes called the descending factorial, falling sequential product, or lower factorial) is defined as the polynomial [Steffensen p.8]

$$(x)_n = x(x - 1)(x - 2)\dots(x - n + 1) = \prod_{k=1}^n (x - k + 1) = \prod_{k=0}^{n-1} (x - k) \tag{1.11}$$

Gauss' Relations for Contiguous Functions:

The six functions $F(a \pm 1, b; c; z), F(a, b \pm 1; c; z), F(a, b; c \pm 1; z)$ are called contiguous to $F(a, b; c; z)$. Relation between $F(a, b; c; z)$ and any two contiguous functions have been given by Gauss [Abramowitz p.558(15.2.19)]. The contiguous relation which is used in this paper is defined as

$$b {}_2F_1 \left[\begin{matrix} a, & b + 1 \\ c & \end{matrix} ; z \right] = (b - c + 1) {}_2F_1 \left[\begin{matrix} a, & b \\ c & \end{matrix} ; z \right] + (c - 1) {}_2F_1 \left[\begin{matrix} a, & b \\ c - 1 & \end{matrix} ; z \right] \tag{1.12}$$

2. Main Generalized Formula

$${}_2F_1 \left[\begin{matrix} a, & b \\ \frac{a+b-n}{2} & \end{matrix} ; \frac{1}{2} \right] =$$

$$\begin{aligned}
 &= \sum_{k=0}^{\infty} \left[\frac{\left\{ 2^{a+b-k+\frac{(-a-b+n)}{2}} \Gamma\left(\frac{a+b-n}{2}\right) \Gamma\left(a+b+\frac{(-a-b+n)}{2}\right) (-a+\frac{a+b-n}{2})_k (-b+\frac{a+b-n}{2})_k \right\}}{k! \Gamma(a) \Gamma(b) (1-a-b+\frac{a+b-n}{2})_k} + \right. \\
 &\quad \left. + \frac{\left\{ 2^{-k} \Gamma(-a-b+\frac{a+b-n}{2}) \Gamma\left(\frac{a+b-n}{2}\right) (a)_k (b)_k \right\}}{k! \Gamma(-a+\frac{a+b-n}{2}) \Gamma(-b+\frac{a+b-n}{2}) (1+a+b+\frac{(-a-b+n)}{2})_k} \right] \\
 &\quad \text{for } \frac{(-a-b-n)}{2} \notin z
 \end{aligned} \tag{2.1}$$

3. Integral Representation of Main Generalized Formula

$${}_2F_1 \left[\begin{matrix} a, b & ; & \frac{1}{2} \\ \frac{a+b-n}{2} & ; & \frac{1}{2} \end{matrix} \right] = \frac{1}{\Gamma(b)} \int_0^{\infty} e^{-t} t^{-1+b} {}_1F_1 \left[a; \frac{a+b-n}{2}; \frac{t}{2} \right] dt \text{ for } \text{Re}(b) > 0. \tag{3.1}$$

$$\begin{aligned}
 &= -\frac{\iota \Gamma\left(\frac{a+b-n}{2}\right)}{2\pi \Gamma(a) \Gamma(b) \Gamma\left(\frac{a-b-n}{2}\right) \Gamma\left(\frac{-a+b-n}{2}\right)} \int_{-\iota\infty+\gamma}^{\iota\infty+\gamma} 2^s \Gamma(a-s) \Gamma(b-s) \Gamma(s) \Gamma\left(\frac{-a-b-n+2s}{2}\right) ds \\
 &\quad \text{for } \max\left(0, \frac{1}{2} \text{Re}(a+b+n)\right) < \gamma < \min(\text{Re}(a), \text{Re}(b))
 \end{aligned} \tag{3.2}$$

4. Special cases of the formula

If $a = b = 1$, then

$$\frac{d}{dn} \left[{}_2F_1 \left[\begin{matrix} 1, 1 & ; & \frac{1}{2} \\ \frac{2-n}{2} & ; & \frac{1}{2} \end{matrix} \right] \right] = \frac{1}{8} \left[-4\psi^{(0)}\left(\frac{1}{2} - \frac{n}{4}\right) + 4\psi^{(0)}\left(-\frac{n}{4}\right) + n \left\{ \psi^{(1)}\left(\frac{1}{2} - \frac{n}{4}\right) - \psi^{(1)}\left(-\frac{n}{4}\right) \right\} \right] \tag{4.1}$$

$$\int {}_2F_1 \left[\begin{matrix} 1, 1 & ; & \frac{1}{2} \\ \frac{2-n}{2} & ; & \frac{1}{2} \end{matrix} \right] dn = \left[2n \log \Gamma\left(\frac{1}{2} - \frac{n}{4}\right) - 2n \log \Gamma\left(-\frac{n}{4}\right) + 8\psi^{(-2)}\left(\frac{1}{2} - \frac{n}{4}\right) - 8\psi^{(-2)}\left(-\frac{n}{4}\right) \right] + \text{Constant} \tag{4.2}$$

If $a = b = 0$, then

$$\begin{aligned}
 {}_2F_1 \left[\begin{matrix} 0, 0 & ; & \frac{1}{2} \\ \frac{-n}{2} & ; & \frac{1}{2} \end{matrix} \right] &= \sum_{k=0}^{\infty} \frac{{}_2F_1(k, k; k - \frac{n}{2}; z_0) ((0)_k)^2 (\frac{1}{2} - z_0)^k}{k! (-\frac{n}{2})_k} \\
 &\text{for } (\text{not}(z_0 \in R \text{ and } 1 \leq z_0 \leq \infty))
 \end{aligned} \tag{4.3}$$

References

- [1] Abramowitz, Milton., A and Stegun, Irene ; *Handbook of Mathematical Functions with Formulas , Graphs , and Mathematical Tables*, National Bureau of Standards, 1970.
- [2] Aomoto, Kazuhiko, Kita, Michitake ; *Theory of Hypergeometric functions*, Springer Tokyo Dordrecht Heidelberg London New York, 2011.
- [3] Gauss, C. F. ; *Disquisitiones generales circa seriem infinitam ...* , Comm. soc. reg. sci. Gott. rec., 2(1813), 123-162.
- [4] Koepf, W.; *Hypergeometric Summation: An Algorithmic Approach to Summation and Special Function Identities*. Braunschweig, Germany: Vieweg, 1998.
- [5] Luke, Y. L.; *Mathematical functions and their approximations*. Academic Press Inc., London,, 1975.
- [6] P. Appell, Sur une formule de M. Tisserand et sur les fonctions hypergéométriques de deux variables, *J. Math. Pures Appl.*, (3) 10 (1884) 407-428.
- [7] Prudnikov, A.P., Brychkov, Yu. A. and Marichev, O.I.; *Integral and Series Vol 3: More Special Functions*, Nauka, Moscow, 2003.
- [8] Salahuddin, Kholā, R. K.; New hypergeometric summation formulae arising from the summation formulae of Prudnikov, *South Asian Journal of Mathematics*, 4(2014), 192-196.
- [9] Steffensen, J. F.; *Interpolation (2nd ed.)*, Dover Publications, U.S.A, 2006.