

## Bare surfaces analysis within Ograzhden Mountain (Bulgarian part)

Rosen Iliev

Institute for Space Research and Technology, Bulgarian Academy of Sciences, Sofia, Bulgaria

Email address: [ilievrosen@space.bas.bg](mailto:ilievrosen@space.bas.bg)

### Abstract

The presence of bare surfaces is one of the main indicators for the degradation of natural ecosystems. The spread of bare surfaces and respectively the lack of vegetation cover is a serious threat to the sustainable development of complex environments such as mountains. The present study focuses on the spatial-temporal relationships of the bare surfaces with the local environment within the Bulgarian part of the small border mountain of Ograzhden. Based on temporal imageries from the Landsat 8 satellite, an analysis of the spatial distribution and tendencies in the development of bare surfaces has been made over a two-year period of time. Overall, the results of the research show a reduction in the total area of bare surfaces on the one hand and a slow environmental recovery on the other. Unfortunately, the positive changes are not ubiquitous, but describe only certain areas of the territory of interest.

**Keywords:** Ograzhden, mountain, bare surfaces, environment, remote sensing, Bulgarian

### Introduction

Most common indicator for negative development of the natural environments is the presence of bare surfaces. In general, bare surfaces represent territories on Earth's surface not covered by vegetation. The term combines all territories occupied by bare soils and rock outcrops. Unfortunately it is one of the most important and typical land cover types all over the world. Bare land, whether seasonal or permanently bare, plays an important role in many ecosystems [1].

Generally, in the temperate regions the rapid development of this type of land cover often is a result of an adverse human impact on the environment, such as mineral exploration, deforestation, intensive cattle breeding and agriculture, etc.

Nowadays, remote sensing is one of the advanced tools for providing information of the spatial-temporal changes of the areas vulnerable to degradation [2,3], especially the mountain environments [4]. Multitemporal data obtained from satellite images provide good geographical information for effectively managing of environmental processes within the mountain regions. Due to the difficult access to most mountain regions – difficult for physical and/or political reasons – remote sensing is often the only way for investigating large sections of the Earth's surface [5]. One of the most common applications of remote sensing is mapping and characterizing the surface cover. Manual and semi-automatic segmentation of optical images for vegetation, water, snow, rock, human objects, etc. can be based on panchromatic or color images. Multi-spectral remote sensing offers the opportunity for automatic classification of surface cover utilizing the variation in reflectivity with wavelength, which differs for most surface types [5, 6].

The main goal of this study is to describe the vegetation cover spread and its relation to bare surfaces within the Bulgarian part of Ograzhden Mountain. The research investigates the spatial-temporal peculiarities of the local environment using data from Landsat 8 satellite for a period of two years (08.08.2017 - 12.08.2019). This is a quantitative and qualitative assessment of the status and spatial distribution of the bare surfaces and vegetation cover. The results of the study show positive trends, though not everywhere.

## Study Area, Data and Working Methods

### Study area

The object of present study is Bulgarian part of the small border mountain of Ograzhden (Fig.1) (41°57' - 41°39' N; 22°95' - 23°25' E). It is integral part of the Osogovo-Belasitsa mountain range and currently it is shared between Republik of Northern Macedonia (about 54%) and Republik of Bulgaria (about 46%). The total area of the Bulgarian part is 331 km<sup>2</sup> and has a length of 24 km and maximum width of 18 km. To the north it borders Maleshevo Mountain, to the south is the valley of the Strumeshnitsa River, which separates it from Belasitsa Mountain. To the east, the Struma River Valley separates it from Pirin Mountain. The western border coincides with the state border with Republik of Northern Macedonia. The highest point of Ograzhden Mountain - the Ograzhdanets peak (1748 m), is located on Northern Macedonian territory. Within the Bulgarian part, the highest point is Bilska chuka peak (1644 m), which is an important hydrographic knot. Other more well-known peaks are Markovi kladentsi peak (1523 m), Muratov peak (1398 m), Kukovski chukar peak (1233 m) [7], etc.




Fig.1: Geographical location and 3D view of the study area (Base map: Google Terrain).

## Image Processing Method

Image processing methods transform the multispectral satellite data into images that enhance study features in contrast with the background [8]. Image fusion techniques deal with integration of complementary and redundant information from multiple images to create a composite image that contains a better description of the scene [9]. In the present study image processing techniques such as Band composite, Grid clustering and Change detection are applied. The band combination 5 (for Red) (wavelength 0,85-0,88  $\mu\text{m}$ ), 6 (for Green) (wavelength 1,57-1,65  $\mu\text{m}$ ) and 7 (for Blue) (wavelength 2,11-2,29  $\mu\text{m}$ ) allows differentiation of bare surfaces from vegetation. In the attached image (Fig.2), the bare surfaces are presented in a light blue colour and the vegetation in an orange colour. However, it is difficult to visually differentiate natural bare surfaces and the built-up areas in one image due to their high complexity and similarity of spectral response patterns, especially in a mixture of pixels with heterogeneous objects [10]. Within the surveyed area however, the settlements are very small and with a steadily decreasing population, so their total share of "bare surfaces" is minimal.

The vegetation and bare surfaces are extracted through unsupervised k-means clustering of grids, based on Hill-Climbing method [11]. The vegetation cover additionally was categorized into two classes – light (grass and individual shrubs) and heavy (closed forest and dense shrub vegetation) ones. Then, using change detection technique the spatial-temporal peculiarities of the extracted classes was analyzed.




Fig.2: Composite image (bands 5, 6 and 7) of the study area. The bare surfaces are colored in light blue and vegetation is colored in orange.

## Data and Software

The imagery data for the present study were acquired from Landsat 8 satellite using the United States Geological Survey (USGS) online interface, Earth Explorer (online available at <https://earthexplorer.usgs.gov/>). The imageries are available on a Universal Transverse Mercator (UTM) mapping grid at 30x30 m pixel size. The satellite data were acquired as zipped Georeferenced Tagged Image File Format (GeoTIFF) representing systematically terrain corrected data (L1T) for period of time 08.08.2017 and 12.08.2019.

The software used for processing and information extraction of imagery data are ArcGIS 10.1 and SAGA-GIS [12].

### Results and Discussion

The results of the environmental monitoring of the bare surfaces within the study area for the period of time 08.08.2017 - 12.08.2019 show a reduction in their total area by 5% (Table 1). On the other hand, there is a restoration of the vegetation cover as a whole (+3%), especially the heavy one (+10%) (Table 2).

Table 1: Vegetation cover and bare surfaces distribution within the study area for the period of time 08.08.2017 – 12.08.2019

Land Cover	August 2017			August 2019		
	Area (km <sup>2</sup> )	Area (ha)	Relative share (%)	Area (km <sup>2</sup> )	Area (ha)	Relative share (%)
<b>Vegetation</b>	<b>239</b>	<b>23900</b>	<b>72,2</b>	<b>245,6</b>	<b>24566</b>	<b>74,4</b>
<i>Heavy</i>	<i>136</i>	<i>13600</i>	<i>41,1</i>	<i>150,4</i>	<i>15044</i>	<i>45,5</i>
<i>Light</i>	<i>103</i>	<i>10300</i>	<i>31,1</i>	<i>95,2</i>	<i>9522</i>	<i>28,9</i>
<b>Bare surface</b>	<b>92</b>	<b>9203</b>	<b>27,8</b>	<b>85,4</b>	<b>8538</b>	<b>25,6</b>

Table 2: Vegetation cover and bare surfaces spatial changes within the study area for the period of time 08.08.2017 – 12.08.2019

Land Cover transition	Area (km <sup>2</sup> )	Area (ha)
Heavy vegetation >> Light vegetation	2,29	229
Bare surface >> Light vegetation	10,12	1012
Light vegetation >> Bare surface	3,48	348
Light vegetation >> Heavy vegetation	16,69	1669

In spatial terms, positive changes are not ubiquitous, but there is a rigorous geographical determination (Fig.3). Restoration of the vegetation cover is mainly related to the western parts of the study area. To the east, the situation is still unfavorable. Here the erosion processes are highly developed and give prerequisites for the occurrence of adverse hydro-climatic phenomena.

The intensive depopulation processes within the study area and the disappearance of cattle breeding in the last decades have contributed significantly to the positive trend. Unfortunately, in the eastern parts of the study area, the natural environment is so badly damaged thus special measures are needed to overcome the negative situation.




Fig.3: Spatial distribution of the vegetation cover and bare surfaces within the study area. From attached maps are visible the reduction of the total area of bare surfaces and an increase of vegetation cover. Positive changes are geographically related to the western parts of the study area, while to the east the situation is still very worse.

## Conclusions

In the present study an analysis of the condition of bare surfaces and vegetation cover within the Bulgarian part of Ograzhden Mountain was made. Environmental monitoring using remote sensing has shown positive trends and reduced areas of bare surfaces. This fact, along with the increase in vegetation cover on the other hand, is an indicator of environmental restoration. The vegetation cover and especially the forests are the main natural resource of mountain ecosystems such as Ograzhden Mountain. Its slow restoration in the medium to long term would reduce the negative impact of erosion processes (and probably some human influence) on these lands and guarantee the sustainable development of the local ecosystem.

## References

1. Rasul, A., Balzter, H., Faqe Ibrahim, G., Hameed, H., Wheeler, J., Adamu, B. and Najmaddin, P. (2018) Applying Built-Up and Bare-Soil indices from Landsat 8 to cities in dry climates. *Land*, 7, 81, 13 p.; doi:10.3390/land7030081
2. Liang, D., Zuo, Y., Huang, L., Zhao, J., Teng, L. and Yang, F. (2015) Evaluation of the consistency of MODIS land cover product (MCD12Q1) based on Chinese 30 m globeland 30 datasets: A case study in Anhui Province, China. *ISPRS Int. J. Geo-Inf.*, 4, 2519–2541.
3. Satyanarayana, B., Thierry, B., Seen, D.L., Raman, A.V. and Muthusankar, G. (2001) Remote sensing in mangrove research-relationship between vegetation indices and dendrometric parameters: A case for Coringa, east coast of India. In proceeding of the 22nd Asian Conference on Remote Sensing, Singapore, 5–9 November 2001; 5, 9 p.
4. Mohajane, M., Essahlaoui, A., Oudija, F., El Hafyani, M., El Hmaid, A., El Ouali, A., Randazzo, G. and Teodoro, A. (2018) Land Use/Land Cover (LULC) using Landsat data series (MSS, TM, ETM+ and OLI) in Azrou Forest, in the Central Middle Atlas of Morocco. *Environments*, 5, 131, 16 p.; doi:10.3390/environments5120131
5. Kaab, A. (2005) Remote Sensing of Mountain Environments. UNESCO publications: Proceedings Second and Third GLOCHAMORE Workshop, 92-99.
6. Campbell, J.B. (2002) Introduction to Remote Sensing. Third edition, London: Taylor and Francis.
7. Nikolov, V., Yordanova, M. and Boteva, I. (2013) Mountains in Bulgaria. Third edition, Sofia: "Prof. Marin Drinov" Academic Press (in Bulgarian)
8. Da Cunha Frutuoso, R.M. (2015) Mapping hydrothermal gold mineralization using Landsat 8 data. A case study in Chaves license, Portugal. Porto: University of Porto, 85p.
9. Wen, C.Y. and Chen, J.K. (2004) Multi-resolution image fusion technique and its application in forensic science *Forensic Science International*, 140, 217-232.
10. He, C., Shi, P., Xie, D., and Zhao, Y. (2010) Improving the normalized difference built-up index to map urban built-up areas using a semiautomatic segmentation approach. *Remote Sensing Letters*, 1, 213–221.
11. Rubin, J. (1967) Optimal Classification into Groups: An Approach for Solving the Taxonomy Problem. *Journal of Theoretical Biology*, 15, 103-144.
12. Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., Wehberg, J., Wichmann, V. and Boehner, J. (2015) System for Automated Geoscientific Analyses (SAGA) v. 2.1.4. *Geosci. Model Dev.*, 8, 1991-2007, doi:10.5194/gmd-8-1991-201